

Affine Anosov diffeomorphisms of affine manifolds.

by Aristide Tsemo
The Abdus Salam International Centre for Theoretical Physics
Strada Costiera, 11
Trieste Italy.
Email tsemo@ictp.trieste.it

Abstract.

In this paper, we show that a compact affine manifold endowed with an affine Anosov transformation map is finitely covered by a complete affine nilmanifold. This answers a conjecture of Franks for affine manifolds.

0. Introduction.

An n -affine manifold (M, ∇_M) is an n -differentiable manifold M endowed with a locally flat connection ∇_M , that is a connection ∇_M whose curvature and torsion forms vanish identically. The connection ∇_M defines on M an atlas (affine) whose transition functions are affine maps. The connection ∇_M pulls-back on the universal cover $\hat{M}$ of M , and defines on it a locally flat connection $\nabla_{\hat{M}}$. The affine structure of $\hat{M}$ is defined by a local diffeomorphism $D_M : \hat{M} \rightarrow \mathbb{R}^n$ called the developing map. The developing map gives rise to a representation $h_M : \pi_1(M) \rightarrow \text{Aff}(\mathbb{R}^n)$ called the holonomy. The linear part $L(h_M)$ of h_M is the linear holonomy. The affine manifold (M, ∇_M) is complete if and only if D_M is a diffeomorphism. This means also that the connection ∇_M is complete.

A diffeomorphism f of M is an Anosov diffeomorphism if and only if there is a norm $\| \cdot \|$ associated to a riemannian metric $<, >$, a real number $0 < \lambda < 1$ such that the tangent bundle TM of M is the direct summand of two bundles TM^s and TM^u called respectively the stable and the unstable bundle such that:

$$\| df^m(v) \| \leq c\lambda^m \| v \|, v \in TM^s$$

$$\| df^m(w) \| \geq c\lambda^{-m} \| w \|, w \in TM^u;$$

where d is the usual differential, c a positive real number, and m a positive integer.

The stable distribution TM^s (resp. the unstable distribution TM^u) is tangent to a topological foliation $\mathcal{F}^s$ (resp. $\mathcal{F}^u$).

The property for a diffeomorphism to be Anosov is independent of the choice of the riemannian metric if M is compact. In this case we can suppose that c is 1.

A Franks's conjecture asserts that an Anosov diffeomorphism f of a compact manifold M is C^0 -conjugated to an hyperbolic infranilautomorphism. This conjecture is proved in [1] with the assumptions that f is topologically transitive

it's stable and unstable foliations are C^∞ , it preserves a symplectic form or a connection.

The goal of this paper is to characterize compact affine manifolds endowed with affine Anosov transformations. More precisely, we show:

Theorem 1. *Let (M, ∇_M) be a compact affine manifold, and f an affine Anosov transformation of M , then (M, ∇_M) is finitely covered by a complete affine nilmanifold.*

1. The Proof of the main theorem.

The main goal of this part is to show theorem 1. In the sequel, (M, ∇_M) will be an n -compact affine manifold endowed with an affine Anosov diffeomorphism f . The stable foliation $\mathcal{F}^s$ (resp. the unstable foliation $\mathcal{F}^u$) pulls-back on the universal cover $\hat{M}$ to a foliation $\hat{\mathcal{F}}^s$, (resp. $\hat{\mathcal{F}}^u$).

Let (U, ϕ) be an affine chart of M , the restriction of a riemannian metric of M to U will be said an euclidean metric adapted to the affine structure of U if its restriction to U is the pulls-back by ϕ on U , of the restriction to $\phi(U)$ of an euclidean metric of $\mathbb{R}^n$.

Proposition 1.1. *The stable and the unstable distributions of f define on (M, ∇_M) foliations whose leaves are immersed affine submanifolds.*

Proof. Let x be an element of M , $\| \cdot \|$ a norm associated to a riemannian metric $\langle, \rangle$ of M . Let v be an element of $T_x M^s$ the subspace of $T_x M$ tangent to $\mathcal{F}^s$, we have $\| df^m(v) \| \leq \lambda^m \| v \|$ where $0 < \lambda < 1$, and $m \in \mathbb{N}$. Let (U, ϕ) be an affine chart which contains a point of accumulation of the sequence $(f^p(x))_{p \in \mathbb{N}}$. We can suppose that the restriction of $\langle, \rangle$ to U is euclidean and adapted to the affine structure. Let $p > p'$ such that $f^p(x)$ and $f^{p'}(x)$ are elements of U , and $v \in T_{f^{p'}(x)} M^s$, we have $\| df^{p-p'}(v) \| \leq \lambda^{p-p'} \| v \|$. This implies that for every element $y = f^{p'}(x) + w$, $w \in T_{f^{p'}(x)} M^s$ and $f^{p-p'}(y) \in U$ is an element of the stable leaf of $f^{p'}(x)$ since the distance between $f^{n_q}(x)$ and $f^{n_q}(y)$ goes to zero for a subsequence $n_q > p$ such that $f^{n_q}(x)$ is an element of U . We deduce that this leaf is an immersed submanifold. The result for the unstable foliation is deduced considering f^{-1} .

Proposition 1.2. *The leaves of $\mathcal{F}^s$ are geodesically complete for the affine structure.*

Proof. Let x be an element of M , as M is supposed to be compact, the sequence $(f^m(x))_{m \in \mathbb{N}}$ has a point of accumulation y . Let U_y be an open set containing y such that there is a strictly positive number r , such that for every $z \in U_y$, $v \in T_z M$ whose norm is less than r for a given riemannian metric, a (affine) geodesic from x whose derivative at 0 is v is defined at 1. Let w be an element of $T_x M^s$ such that $\| df^p(w) \| \leq r$. Without loss generality, we can suppose that $f^p(x) \in U_y$, the (affine) geodesic from $f^p(x)$ whose derivative

is $df^p(w)$ at 0 is defined at 1. This implies that the geodesic from x which derivative at 0 is w is defined at 1, since f^{-1} is an affine map. This shows the result.

It is a well-known fact that an Anosov's diffeomorphism of a compact manifold as a periodic fixed point. We will assume up to change f by an iterated that f has a fixed point x . There exists also a map $F : \hat{M} \rightarrow \hat{M}$ over f which fixed the element $\hat{x}$ over x .

Proposition 1.3. *Let $\hat{y}$ and $\hat{z}$ be two elements of $\hat{\mathcal{F}}_t^u$, where t is an element of $\hat{M}$, then the images of $\hat{\mathcal{F}}_{\hat{t}}^s$ and $\hat{\mathcal{F}}_{\hat{z}}^s$ by the developing map are parallel affine subspaces.*

Proof. Let $\hat{y}$ be an element of $\hat{\mathcal{F}}_t^u$, it is sufficient to show that $D(\hat{\mathcal{F}}_t^s)$ and $D(\hat{\mathcal{F}}_{\hat{y}}^s)$ are parallel affine subspaces.

We know that the tangent bundle of a simply connected affine manifold is trivial. We have $T\hat{M} = \hat{M} \times T_t\hat{M}$. Let $<, >'$ be a riemannian metric on M which pulls back on $\hat{M}$ is $<, >$. The map F is Anosov relatively to $<, >$. Without loss generality, we can assume that the distributions tangent to $\hat{\mathcal{F}}^s$ and $\hat{\mathcal{F}}^u$ are orthogonal.

Let w be a vector of $T\hat{M}_{\hat{y}}$ tangent to $\hat{\mathcal{F}}_{\hat{y}}^s$. We can write $w = s + u$ where s is a vector of $T\hat{M}_{\hat{t}}$ tangent to $\hat{\mathcal{F}}^s$ and u is a vector of $T\hat{M}_{\hat{t}}$ tangent to $\hat{\mathcal{F}}^u$. The vector u is also an element of $T\hat{M}_{\hat{y}}$ tangent to $\hat{\mathcal{F}}_{\hat{y}}^u$ since the unstable foliation is affine (we can identify the projection on the second factor of $T\hat{M} \times T\hat{M}_{\hat{t}}$ of vectors tangent to $\hat{t}$ and $\hat{y}$ since $T\hat{M}$ is trivial). This implies that $\|dF_{\hat{y}}^n(u)\| \geq \lambda^{-n} \|u\|$ for $0 < \lambda < 1$. But on the other hand we have $\|dF_{\hat{y}}^n(s + u)\| \leq \lambda^n \|(s + u)\|$, which implies that the limit of $\|dF_{\hat{y}}^n(s + u)\|$ is zero when n goes to infinity. We deduce that $u = 0$ since we have supposed that the distributions tangent to $\hat{\mathcal{F}}^s$ and $\hat{\mathcal{F}}^u$ are orthogonal, so $\|dF_{\hat{y}}(s + u)\| = \|dF_{\hat{y}}(s)\| + \|dF_{\hat{y}}(u)\|$. This implies the result.

The images by D_M of the leaf $\hat{\mathcal{F}}_{\hat{t}}^s$ of $\hat{\mathcal{F}}^s$ and $\hat{\mathcal{F}}_{\hat{t}}^u$ of $\hat{\mathcal{F}}^u$ are affine subspaces of $\mathbb{R}^n$ whose direction are direct summand of $\mathbb{R}^n$. Since for every element z of $\hat{\mathcal{F}}_{\hat{t}}^s$ the leaf of $\mathcal{F}^u$ passing by z is complete, we deduce that the developing map is surjective.

Proposition 1.4. *The affine manifold (M, ∇_M) is complete.*

Proof. Let $\hat{t}$ be an element of $\hat{M}$, and $\hat{E}_{\hat{t}}$ the set of elements y of $\hat{M}$ such that there is an element z in $\hat{\mathcal{F}}_{\hat{t}}^u$ such that y is an element of $\hat{\mathcal{F}}_z^s$. The image of $\hat{E}_{\hat{t}}$ by D_M is $\mathbb{R}^n$, and the restriction of D_M to $\hat{E}_{\hat{t}}$ is injective. The set $\{\hat{E}_{\hat{t}}, \hat{t} \in \hat{M}\}$ is a partition of $\hat{M}$ by disjoint open sets. It has only one element since $\hat{M}$ is connected. We deduce that (M, ∇_M) is complete.

Remark.

The existence of an affine Anosov transformation $\hat{f}$ on the universal cover of a compact affine manifold (M, ∇_M) which pulls forward onto a diffeomorphism f of M does not implies that f is an Anosov diffeomorphism as shows the following example:

Let $Hopf(n)$ be the quotient of $\mathbb{R}^n/\{0\}$ by an homothetic h_λ whose ratio λ is such that $0 < \lambda < 1$. It is a compact affine manifold. Every homothetic h_c which ratio c is positive and different from 1 and λ is an Anosov diffeomorphism of $\mathbb{R}^n$ endowed with an euclidean metric. But the pulls forward of h_c is an isometry of $Hopf(n)$ endowed with the pulls forward of the riemannian metric of $\mathbb{R}^n/\{0\}$ defined as follows:

$$\frac{1}{\|x\|^2} < u, v >$$

where x is an element of $\mathbb{R}^n/\{0\}$, and u, v are elements of its tangent space.

Proof of theorem 1.

First we show that the module of the eigenvalues of the elements of the linear holonomy of (M, ∇_M) is 1.

Let (A, a) be an element of $\pi_1(M)$ such that A has an eigenvector u associated to an eigenvalue b (which may be a complex number) whose norm is different from 1.

It is a well-known fact that an Anosov diffeomorphism of a compact manifold has a fixed periodic point. Without loss of generality, we will assume that up to change f by an iterated that f has a fixed point x . This implies that up to a change of coordinates, there exists on $\mathbb{R}^n$ a linear map F over f .

Put $x = p(0)$ where p is the covering map and consider a riemannian metric $<, >'$ on M whose restriction on an affine neighborhood N of x is euclidean adapted to the affine structure. Expressing on N the fact that F is an Anosov diffeomorphism using the metric $<, >'$, one obtains that $\mathbb{R}^n = U \oplus V$, where U and V are two subvectors spaces such that there exists a number $0 < \lambda < 1$ such that

$$\begin{aligned} \|F^n(u)\| &\leq \lambda^n \|u\|, u \in U, \\ \|F^n(v)\| &\geq \lambda^{-n} \|v\|, v \in V \end{aligned}$$

where u , and v are respectively elements of U and V and $\|\cdot\|$ is a norm associated to an euclidean metric $<, >$ of $\mathbb{R}^n$. The subvectors spaces U and V pulls forward respectively onto $T_x M^s$ and $T_x M^u$. We will assume that they are orthogonal with respect to $<, >$. The vectors spaces U and V are stable by the linear holonomy (see proposition 1.3)

Put $u = u_1 + u_2$, where u_1 and u_2 are respectively elements of U' and V' the complexified vectors spaces respectively associated to U and V .

Without restrict the generality, one can assume that after eventually having changed γ by γ^{-1} and (or) f by f^{-1} that u_1 is not zero and that the norm of b is strictly superior to 1.

Let q be a positive integer, consider the smallest positive integer n_q such that $F^{n_q} \circ \gamma^q$ has a fixed point $\hat{m}_q$. The integer n_q exists since A preserves U

and V . Let y be a point of accumulation of the sequence $p(\hat{m}_q)$. Up to replace $(\hat{m}_q)$ by a subsequence and replace $\hat{m}_q$ by another element over m_q , we can suppose that the sequence $(\hat{m}_q)$ converges to $\hat{y}$ over y . The linear part of the elements over f^{n_q} which fixe the elements over m_q have the same eigenvalues since they are conjugated. Consider a riemannian metric of M which pulls back on $\mathbb{R}^n$ coincides on a neighborhood of $\hat{y}$ with an euclidean metric.

1. The sequence (n_q) is bounded.

The restriction of the linear part of the element of f^{n_q} conjugated to $F^{n_q} \circ \gamma^q$ which fixe $\hat{m}_q$ at U is contractant (for the last euclidean metric) with ratio λ^{n_q} ($0 < \lambda < 1$). This is not possible since the restriction of the linear part of this element to U' has the same eigenvalues than the restriction of $F^{n_q} \circ A^q$ to U' , and the limit when q goes to infinity of the norm of $F^{n_q} \circ A^q(u_1)$ for the hermitian metric associated to the last euclidean metric is infinity, since the sequence n_q is bounded and the norm of $b > 1$.

2. The sequence (n_q) is not bounded.

Up to change (n_q) by a subsequence, we can suppose that (n_q) goes to infinity. The linear map $F^{n_q-1} \circ \gamma^q$ does not have a fixed point. Its linear part has the eigenvalue 1 associated to the eigenvector v_q . Write $v_q = v_{1q} + v_{2q}$ where v_{1q} and v_{2q} are respectively element of U and V . If v_{1q} is not zero, then $\|F^{n_q} \circ A^q(v_{1q})\| = \|F(v_{1q})\|$ (the norm considered is the precedent euclidean norm); the restriction of $F^{n_q} \circ A^q$ cannot be contractant with ratio λ^{n_q} for q enough big since the sequence (n_q) goes to infinity.

If v_{2q} is not zero, then $\|F^{n_q} \circ A^q(v_{2q})\| = \|F(v_{2q})\|$, the restriction of $F^{n_q} \circ A^q$ to V cannot be dilatant (for the last euclidean metric) with ratio λ^{-n_q} for q enough big since n_q goes to infinity. There is a contradiction since the eigenvalues of the restriction of $F^{n_q} \circ A^q$ to U' (resp to V') coincide with the eigenvalues of the restriction to U' of the linear part of the conjugated to $F^{n_q} \circ \gamma^q$ which fixes $\hat{m}_q$ (resp. the eigenvalues of the restriction to V' of the linear part of the element conjugated to $F^{n_q} \circ \gamma^q$ which fixes $\hat{m}_q$). We deduce that the eigenvalues of A are roots of unity.

We deduce from the first paragraph of [4] p.6 that $\pi_1(M)$ has a subgroup of finite index G such that the eigenvalues of the linear part of the elements of G are 1. We deduce from [3] that the quotient of $\mathbb{R}^n$ by G is a complete affine nilmanifold.

Remark.

Let (M', ∇') be the finite cover of (M, ∇) which fundamental group is G , and f' the pulls-back of f to M' . The map $H^n(M, \mathbb{R}) \rightarrow H^n(M, \mathbb{R})$, $\alpha \rightarrow (f'^2)^* \alpha$ is the identity since f'^2 is a diffeomorphism which preserves the orientation. Let ω be the parallel form of (M', ∇') , we deduce that (f'^2) preserves the parallel volume form of (M', ∇') ; then a well-known result of Anosov implies that f'^2 is ergodic and its periodic points are dense.

Bibliography.

- [1] Benoist Y., Labourie F., Sur les difféomorphismes d'Anosov affines à feuilles stables et instables différentiables, *Invent. Math.* 111 (1993) 285-308.
- [2] Benoist Y., Foulon P., Labourie F., Flots d'Anosov à distributions stable et instable différentiables, *J. Amer. Math. Soc.* 5 (1992) 33-74.
- [3] Fried D., Goldman, W., Hirsch M., Affine manifolds with nilpotent holonomy, *Comment. Math. helv.* 56 (1981) 487-523.
- [4] Goldman W., Hirsch M., A generalization of Bieberbach's theorem, *Invent. Math.* 65 (1981) 1-11.
- [5] Koszul J., Variétés localement plates et convexité, *Osaka J. Math.* 2 (1965) 285-290.
- [6] Smale S., Differentiable dynamical systems, *Bull. Amer. Math. Soc.* 73 (1967) 747-817.